

volume could be estimated by gully length correctively. Also, correct estimation of gully length would estimate the gully cross section correctly in 35% of the samples. Nachtergaele et al. (2001b) in their research in Portugal found a power relationship between gully volume and gully length (equation 4).

$$V = 0.05L^{1.27} \quad (4)$$

In which V is gully volume (m³), L is gully length (m), a and b are 0.05 and 1.27 in Portugal and 0.1839 and -0.2385, respectively in loess plateau In China. Capra et al. (2005) in the model calibration for estimation of gully volume in Sicily, Italia, found the relationship number 5 between gully volume and length. To find higher R square, they used stepwise method in SPSS software to indicate the influence of several indices such as , Catchment area, length of catchments, Catchment gradient, Gully length, Gully slope, 24 hours rainfall, third day rainfall, threshold rainfall equal to 37 mm on the volume of gully erosion. The final equation (6) indicated that R square was a little better than equation 5 ($R^2=0.72$) but the most important factor is the length of gully to predict the volume of gully erosion.

$$V = 0.0082L^{1.416} \quad R^2 = 0.64 \quad (5)$$

$$V = 10^{-4.91} L^{1.557} 24 - hRainfall^{1.327} \quad R^2 = 0.72 \quad (6)$$

Zhang et al. (2007) presented a power relationship between the volume and the length of gullies in the cultivated catchments of black soil region , the northeast of China as follows:

$$V = 0.015L^{1.429} \quad (R^2 = 0.67, P < 0.05) \quad (7)$$

3. Methods

This research was carried out in Golestan province, the northeast of IR. Iran. The region with gully erosion area larger than 5 km² was determined using historical and anecdotal evidence and field survey. The climate of gully regions was determined using Demarton's classification. Six representative gullies were selected to measure their morphometric and soil characteristics in each climate zone. Measurements of top and bottom width and depth were done in the headcut, 25, 50 and 75% of gully length from headcuts. Soil samples were taken from those points. View plan, headcut plan and long profiles of the gullies were surveyed using theodolite. The volume of gully erosion was estimated using partial volumes. The partial volume was calculated by multiplying partial area by the interval between sections. The relationship between the volume of gully erosion as dependent variable and morphometric and soil characteristics of gullies as independent variables was considered using stepwise method in SPSS software.

4. Results

Two watersheds from six watersheds experienced gully erosion in Golestan province. Gully erosion occupied more than 600 km². It developed in four climates including mild desertic arid, cold desertic arid, cold semiarid and mild semiarid. Average annual rainfall was between 200 and 471 mm. The length of represented gullies were between 50 and 110 meters. The depth of the gullies was between 0.7 to 9.5 meters. The top width of the gullies varied between 3.7 and 27.5 meters. The bottom width of the gullies was between 1 and 10.5 meters. The volume of gully erosion was calculated between 133 (Alagol region) and 23000 m³ (Maravehtapeh region). The result of this research reveals that a linear and positive relationship exists between the top width and depth of gully erosion although its R^2 is relatively lower (Table 1). Regions with more clay experienced a lower top width and regions with silt more than 60% indicated the largest top width. The relationship between the top width and depth of gullies is significant in the 5% significance level. This is similar to the relationship (2) presented by US-SCS (1966). This relationship implies that gully widening is happening in Golestan province. In other words, the risk of land degradation (rangeland and rain fed farms) is higher in these regions. Also, there is a linear relationship between W/D ratio and gully length (Table 1). It implies that with increasing one unit in the gully length, 0.66 unit of W/D ratio will be increased.

Table 1 Relationship between morphometric parameters of the Golestan gullies

	standard coefficient	R^2	p
$D = 2.146 + 0.173wt$	0.55	0.26	0.05
$wr/D = 2.27 + 0.006L$	0.66	0.41	0.05

The results of statistical analysis indicated that the volume of gully erosion had positive relationship with length, top and bottom width, depth and form factor and negative one with EC, Ca, SAR, Clay, wb/wt and wt/L . The

volume of gully erosion had a significant correlation with length, top and bottom width ($P < 0.01$) and with from factor, w_t/D ($P < 0.05$).

The simplest linear equation existed between the volume and length of representative gullies. The result indicated that 92% of variation in the volume of gully erosion was interpreted by the length of gullies. In addition to gully length, gully depth increased R^2 to 0.97 but with adding gully top width, W_t , to the relationship, R^2 was not increased significantly (Table 2). Considering the power relationship, there exists the following equation as follows.

$$V = 5.64L^{1.24} \quad R^2 = 0.52 \quad P < 0.01 \quad (8)$$

This relationship in regarding R^2 and exponent is very similar with that presented for ephemeral gullies in the croplands of Sicily, Italy. Also the exponent of the power relationship for gullies in Golestan province is very similar to the one presented for ephemeral gullies in Portugal by Nachtergaele et al. (2001b). The difference between α coefficient in Golestan, 5.64, with 0.0082 in gullies from Sicily, Italia, might be related to the different type of gullies. Sicily gullies are ephemeral and small gullies that could be obliterated by normal tillage operations while Golestan gullies were permanent ones and their dimensions were larger than that of the Sicily gullies.

Table 2 Relationship between the volume of gully and it's morphometric characteristics

variable	(β_1)	(β_2)	(β_3)	R^2	P
L	0.932			0.86	0.01
L,D	0.966	0.325		0.97	0.01
L, D, W_t	0.871	0.227	0.159	0.978	0.01

5. Conclusion

In Golestan province, two out of six watersheds experienced gully erosion. Total area that was covered by gully erosion was over 600 km². Gully erosion occurred in four climates including mild desertic arid, cold desertic arid, cold and mild semiarid. The results indicated a linear and positive relationship between the top width and depth of gullies although R^2 was lower. This implies that widening is occurring in gullies of Golestan province. Regions with silt content higher than 60% experienced the highest top width in this part of I.R.Iran. Statistical analysis with stepwise method indicated that variation of the gully length interpreted 92% of the variation in the volume of erosion. R^2 was improved by adding the gully depth but there was no better R^2 with adding top width parameter. We recommend using the length of gully to estimate the volume of gully erosion in this province. This parameter could be especially measured on the historical evidence such as aerial photos.

6. References

- Capra, A., Mazzara, L.M., Scicolone, B., 2005. Application of the EGEM model to predict ephemeral gully erosion in Sicily, Italy. *CATENA*, 59:133-146.
- Cheng, H., Zou, X., Wu, C., Zhang, C., Zheng, Q. and Jiang, Z., 2007. Morphology parameters of ephemeral gully in Characteristics hillslopes on the Loess plateau of China. *Soil and Tillage Research*, 94, 4-14.
- Hughes, A.O., Prosser, I.P., Stevenson, J., Scott, A., Lu, H., Gallant, J. and Moran, C.J., 2001. Gully erosion mapping for the national land and water resources audit. CSIRO Land and Water, Canberra, Technical report 26/01, 19pp.
- Nachtergaele, J. And Poesen J., 2001a. The Value of a physically based model versus an empirical approach in the prediction of ephemeral gully erosion for loess-derived soils. *GEOMORPHOLOGY*, 40:237-252.
- Nachtergaele, J., Poesen, J., Vandekerckove, L., Oostwoud Wijdenes, D., Roxo, M., 2001b. Testing the ephemeral gully erosion model (EGEM) for two Mediterranean environments. *Earth Surface Processes and Land Forms*, 26:17-30.
- Nachtergaele, J. and Poesen, J., 2002. Spatial and temporal variations in resistance of loess-driven soils to ephemeral gully erosion. *European Journal of Soil Science* 53(3): 449-464.
- Poesen, J., Vandaele, K. and Van Wesemael, B., 1998. Gully erosion: importance and model implications. In J. Boardman and D. Favis-Mortlock(eds.), *Modeling soil erosion by water*, Nato series, vol. 155, Springer Verlag Berlin Herdeiberg.
- Poesen, J., Nachtergaele, J., Verstraeten, G. and Valentin, C., 2003. Gully erosion and environmental change: importance and research needs. *Catena* 50:91-133.
- Radoane, M., Ichim, I., Radoane, N. and Surdeanu, V., 1990. Asupra profilului longitudinal si a factorului de forma a ravenelor din podisul moldovei, *Acad. Romana, stud. Cercet. Geogr. Tom. XXXVII*, 67-74.
- Soil Science Society of America, 2001. *Glossary of Soil Science Terms*. Soil Science Society of America. Madison, WI, <http://www.soils.org/sssagloss/>.

- USDA-SCS, 1966, Technical release No. 32(Geology), US Department of Agriculture. Washington. DC, 125-142.
- Vandkerckhove, L., Poesen, J., Oostwoud Wigdenes D., Gyssels G., Beuselinck, L. and Deluna, E., 2000. Characteristics and controlling factors of bank gullies in two semi-arid Mediterranean environments. *Geomorphology*. 3(1-2):37-58.
- Zhang, Y., Wu, Y., Liu, B., Zheng, Q., Yin, J., 2007. Characteristics and factors controlling the development of ephemeral gullies in cultivated catchments of black soil region. Northeast China. *Soil and Tillage Research*. 96: 28-41.